

January 2018

**Socioeconomic determinants and health care utilization  
among elderly people living in Europe: Evidence from  
the Survey of Health, Ageing and Retirement.**

David Cantarero-Prieto\*  
Marta Pascual-Sáez\*  
Javier Lera Torres\*\*

*\*University of Cantabria and GEN*

*\*\*University of Cantabria*



January 2018

**Socioeconomic determinants and health care  
utilization among elderly people living in Europe:  
Evidence from the Survey of Health, Ageing and  
Retirement**

David Cantarero-Prieto\*

Marta Pascual-Sáez\*

Javier Lera Torres\*

*\*Department of Economics. The University of Cantabria*

# **Socioeconomic determinants and health care utilization among elderly people living in Europe: Evidence from the Survey of Health, Ageing and Retirement**

**David Cantarero-Prieto, Marta Pascual-Sáez and Javier Lera Torres**

Email: david.cantarero@unican.es; \_marta.pascual@unican.es; javierlerat94@gmail.com

Department of Economics,

University of Cantabria

Avenue. Los Castros, s/n, Santander CP 39005, Spain.

**Corresponding Author:**

David Cantarero-Prieto

Department of Economics,

University of Cantabria

Avenue. Los Castros, s/n, Santander CP 39005, Spain.

Email: david.cantarero@unican.es

## **ABSTRACT**

This paper examines health care utilization among elderly people in sixteen European countries using the last wave of the Survey of Health, Ageing and Retirement in Europe (SHARE). Negative Binomial regression is conducted to study the main driving factors behind health care utilization (visits to the General Practitioners, GP; Hospital Stays, HS). The empirical results suggest that age, gender, education level, self-assessed health, health limitations and status and other socioeconomic variables are the main driving factors. We also show that socioeconomic variables do not play the same role in every country. From a policy economic approach, we propose important information to the current debates both in the health economics and social welfare literature. Our findings are relevant and have several implications for policy purposes to enhance efficiency, equity and quality of health care that it can be provided.

**JEL Codes:** I10; I18

**Keywords:** Aging; Discrete choice methods; Count Data Methods; Health; Health Care Utilization; SHARE; Europe

# 1. INTRODUCTION

Population aging has become one of the most important topics in our society since it is expected that the ratio of elderly people over all the population rises from 18% to 28% by 2060 [1]. The European Union is concerned about how these projections are going to affect health care services [2, 3] because elderly people are prone to use more the health care system [4-5]

Therefore, the future of the European health care services depends on national provision of care and the ways that they are organized and financed [6-10]. These health care systems (National Health Services or Social Security Systems) have a great relevance since there is some evidence that the use of health care services differs between public and private sector [11]. Besides, depending on the country, a prescription by the GP is needed to visit a specialist [12] and this waiting time can increase the use of emergency services. These differences can be related with income inequalities [13], which is an important factor determining the health care services utilization. A low-income household must wait till the public system can provide them an appointment but a high-income one can use the private sector and get a faster health care service [14]. The economic status is also a factor behind some mental diseases [15] like dementia or depression which are proved to be the most typical conditions affecting elderly people [16-20].

Hence, the purpose of this study is to examine the main drivers of health care utilization measured by the visits to the General Practitioners (GP) and the Hospital Stays (HS). Once an analysis for the European Union as a whole is done, we focus on the 16 member states which we have data for based on a multinational Survey, the last wave of the Survey of Health, Ageing and Retirement in Europe (SHARE), which provides comparable Gross National individual data. The SHARE sample is nationally representative of individuals who are 50 years old and over. This is an important fact since that using SHARE it is possible to disentangle whether all Member States follow a European pattern in the use of healthcare system.

Our main contributions are as follows. Firstly, the drivers of the visits to the GP and the hospital stays are based on the last wave (6<sup>th</sup>wave) of the SHARE. It does not only include an analysis for each Member State because we also treat the European Union as a whole. Secondly, we analyze if there is a similar pattern in the European Union Member States, proposing that similar policy recommendations for the European Union Member States is not always a good idea (One-size-fits-all policy)

The paper is organized as follows. The second section is based on the theoretical framework, methodology and data where our topic is analyzed. The third section is focused on the empirical analysis and a discussion of the results can be found. And finally, the final one presents the conclusions and policy implications.

## 2. METHODOLOGY AND DATA

### Methods

Count data methods are often used to explain the healthcare utilization [21-24] and it is useful for cases like the European health care services (Table 1). Our aim is explaining the variables *GP* (number of visits to the General Practitioner) and *HS* (hospital stays understood as number of nights a person is hospitalized). The estimation is based on two different distributions that are both estimated by maximum likelihood: Poisson and Negative Binomial. Since Poisson models have the property of equidistribution and has heterokedasticity a parameter capable to capture the randomness is introduced. The binomial negative function of density is the following:

$$\Pr(Y_i = y_i | x_i) = \frac{\Gamma(y_i + v_i)}{\Gamma(y_i + 1)\Gamma(v_i)} \left(\frac{v_i}{v_i + \mu_i}\right)^{v_i} \left(\frac{\mu_i}{v_i + \mu_i}\right)^{y_i} \quad (1)$$

Where

$$\mu_i = E(Y_i | x_i) = \exp(x_i \beta) \quad (2)$$

$$v_i = \frac{1}{\alpha} \mu_i^t \quad t = 0, 1 \quad (3)$$

Depending on the way we define the variance there are two possibilities:

$$v = (1/\alpha) \quad (4)$$

Then

$$E(Y_i | x_i) = \exp(x_i \beta) \quad (5)$$

$$\text{Var}(Y_i | x_i) = (1 + \alpha) \exp(x_i \beta) \quad (6)$$

Or

$$\text{Var}(Y_i | x_i) = \exp(x_i \beta) (1 + \alpha \exp(x_i \beta)) \quad (7)$$

If the parameter  $\alpha$  tends to 0, then it would be a Poisson distribution. It can be tested with a equidispersion contrast ( $H_0: \alpha = 0$ ). If  $\alpha > 0$ , then overdispersion and infra dispersion if  $\alpha < 0$

Zero-inflated model are used for *NH*. The weight given to fact that the probability the variable equals zero is higher in comparison to the other specifications [25] and it is assumed that (8) not generated by the same process than the other observations [24]. It can be specified as follows:

$$\begin{aligned} P(Y_i = 0) &= f_1(0) \\ P(Y_i = y_i) &= (1 - f_1(0)) \frac{f_2(y_i)}{1 - f_2(0)} \quad y_i = 1, 2, \dots \end{aligned} \quad (8)$$

Where  $f_1(0)$  and  $f_2(0)$  are probability functions and  $1 - f_2(0)$  is used to truncate on zero.

## *Data*

Our data came from the 6<sup>th</sup> wave of the Survey of Health, Aging and Retirement in Europe (SHARE) [26,27]. One of the novelties of this new wave is that a new country has been included: Croatia. The SHARE survey is the result of the aim of the European institutions to have a deeper and strongest cooperation with Member States to have good data on elderly people. It covers data from health to socioeconomic status passing through lifestyle which will help us to include more determinants of the health care utilization system. In the Table 2, there is a summary of the most important statistics for each relevant variable. Our dependent variables are based on two health care indicators (*GP* and *HS*).

Moreover, since age has a major impact on the health care utilization and it is different among different groups of age, for the EU we have distinguished between four groups: 50-65, 65-75, 75-85 and +85. In the case of each European country, we include the age as an independent variable with the addition of another variable *AGE2* that is the squared of *Age* in order to consider the quadratic relationship of this variable along lifecycle with our dependent variables.

As previous studies [12, 16, 22], we include gender (*Female*), marital status (*Single*), participation in the labor market (*InLabFor*) and the level of education (*PriEduc*, *SecEduc* and *TerEduc*), as relevant variable although this last one could be correlated with the socioeconomic level of individuals. Because of among the European Union there are different health care systems, we include a dummy variable related with a supplementary insurance (*SupInsu*) could be interesting to understand whether having it or not would affect to the use of the public service.

Multimorbidity has been captured through some proxies: the Number of Chronic Diseases a person has (*NCD*), and the number of limitations in Activities of Daily Living a person has (*ADL*). The self-assessed health has been included in the case of being with good health (*SPGH*). Physical characteristics determine the physical health status and we have included four dummies: Underweight (*UW*), NormalWeight (*NW*), Overweight (*OW*) and Obese (*Obese*). At this regard, we use the Body Mass Index (BMI) that is included in the SHARE survey and then people were classified depending whether their BMI was under 18.5 (*UW*), between 18.5 and 25 (*NW*), between 25 and 30 (*OW*) and more of 30 that corresponds to people suffering from obesity [28]. Hence, the importance of obesity has a double side: first, the direct effect that the illness has and the second is the propensity that people included in this condition could suffer from diabetes or other sicknesses which can make citizens request to have a higher coverage that includes specialists like dieticians or psychologists. [29-32]. Finally, Lifestyle is also important in the predisposition of getting ill and in this case the proxy variable selected was related to smoking (*Smoke*).

### 3. EMPIRICAL RESULTS

#### Visits to General Practitioner (*GP*)

In this section, we compare our findings with those concerning with health care utilization using count data models. Our estimates show in Tables 3 and 4 that , being a female is related to more possibilities of going to the GP than men since they are more likely to be ill from disabling conditions [33]. This gender gap decreases along our sample and turns negative for the group of people of more than 85 years old. It could be possible since women have a longer life expectancy than men, 83.3 and 77.9 respectively [34]. It means that those men that are older than 85 have subjective health that it continues to be substantially worse than women which leads them to visit more to the GP as previous studies have demonstrated [35,36].

Being single has different effects along the cohorts. It produces more GP visits in those people aged 50-65 and +85 and less in the other cases. Marital status is not relevant in any country (excepting for Croatia and Czech Republic) and it does not have a similar slope because in some countries it is negative and in others it is positive probably due to the heterogeneity in cultures

Moreover, active people are less prone to visit GPs and this effect increases when people get older. This result is consistent with some previous literature where it was proved that unemployed people were more likely to use the health care services [37]. The fact of having a supplementary insurance is related with more visits to the GPs and education variables also are important.

Self-assessed health, number of chronic diseases and limitations of daily living are statistically significant in all the elderly groups which mean that they are main driving factors of health care utilization. It is important to notice that most of their values get reduced when changing the cohort but continue having the same relevance. More chronic diseases and limitations increase the probability of visit the GP and having a good self-assessed health has the opposite effect. Besides, when *smoke* is significant, we can observe that it has a positive slope which is consistent with the previous literature. The same can be argued to those people that do not have a normal BMI. Generally, there are more likely to visit the GP than those with a normal weight.

#### Hospital Stays (*HS*)

There are wide variations across European countries in Hospital Stays differences as it shows Tables 5-6. Being female is related to have less hospital stays than men and being single, although is only statistically relevant in the general European Union analysis increases the probability of having more hospital stays. As in the case for *GP* visits, age is the main driving factor of health care utilization but here we observe that...

Being in the labor force reduces, in general, the probability of being hospitalized. For the group of people aged between 75 and 85 it is not a relevant factor, something that can be explained because there are not so many people working at that age, and those who continue active in the labor market may have a fantastic health status.

The health care system (National Health Services or Social Security System) could be also behind the differences in the effect of having a supplementary insurance and its influence over health care utilization. Hence, poor people cannot get a supplementary insurance as the people with a higher socioeconomic status. Related with the proxies of the socioeconomic status, we can point out that those people with tertiary education have less probability of having hospital stays than those with a lower education. Being hospitalized can be consequence of not have treated correctly an illness.

Finally, health status measured by multimorbidity and limitations of daily living and the self-assessed health are main driving factors of health care utilization according to our findings which are consistent with previous literature [36-38]. The first two ones have a clear positive effect on the probability of being hospitalized but a good self-assessed health decrease the probability of being hospitalized. Variables related to BMI are not relevant enough although not having a normal BMI increase HS. The same happens with the variable related with smoke proxy because it is only significant in the first group of age and in some countries although their slope is generally positive.

## 4. CONCLUSIONS

The comparison of the elderly people and some socioeconomic variables regarding their health care utilization in Europe is studied here in order to provide new empirical evidence. We demonstrate that the drivers behind the visits to the General Practitioner and Hospital Stays by using the 6<sup>th</sup> wave of the SHARE are quite similar to those described in the previous literature. Our findings point to the fact that there is a similar pattern in the European Union Member States.

Being older increase the health care utilization and being part of the labor force reduces the use of the healthcare system. People with chronic diseases and limitations in their daily activities make them visit more the General Practitioner and to have longer Hospital Stays. But other factors as education, marital status, gender or supplementary insurance can influence health care utilization in order to reduce it.

The results of this research add to our understanding of the behavior about health care utilization of elderly people across Europe. We really think that health campaigns, a better use of digital medicine or to empower patients to take care by themselves should be designed to explain to the people how health care (primary and hospital) should be used. Because of the heterogeneous nature of General Practitioner visits and Hospital Stays in Europe and the increasing pressure of a growing elderly population, more effort in this direction has to be carried out in order to enhance efficiency, equity and quality of health care systems.



## REFERENCES

1. European Commission (2015): “The 2015 aging report – Economic and budgetary projections for the 28 EU Member States (2013-2060)”
2. European Commission (2013): “Investing in health. Commission Staff Working Document. Social Investment Package” Available at:  
[https://ec.europa.eu/health/sites/health/files/strategy/docs/swd\\_investing\\_in\\_health.pdf](https://ec.europa.eu/health/sites/health/files/strategy/docs/swd_investing_in_health.pdf)
3. World Bank (2017): Development Indicators (WDI) Database. Available at:  
<http://databank.worldbank.org/data/home.aspx>
4. Aguado, A; Rodriguez, D; Flor, F; Sicras, A; Ruiz, A; Prados (2011): “Distribución del gasto sanitario en atención sanitaria primaria según edad y sexo: un análisis retrospectivo”, *Atención Primaria* (2012) 44(3):145-152
5. Meijer, C; Wouterse, B; Polder, J; Koopmanschap, M (2013): “The effect of population aging on health expenditure growth: a critical review”, *Eur J Aging* 10:353-361
6. Thomson, S; Foubister, T; Mossialos, E (2009): “Financing health care in the European Union. Challenges and policy responses”, *Observatory Studies Series N° 17*
7. Abásolo, I; Negrín-Hernández; M.A., Pinilla, J (2014): “Utilización y tiempos de espera: dos vertientes inseparables del análisis de la equidad en el acceso al sistema sanitario público”, *Hacienda Pública Española*, 208(1): 11-38.
8. Åhs, A.M.H; Westerling, R (2006): “Health care utilization among persons who are unemployed or outside the labour force”, *Health Policy*, 78(2): 178-193
9. Jeon, B; Noguchi, H; Kwon, S; Ito, T; Tamiya, N. (2017): “Disability, poverty, and role of the basic livelihood security system on health services utilization among the elderly in South Korea”, *Social Science & Medicine*, 178: 175-183.
10. Hosseinichimeh, N; Martin, E and Weinberg, S (2016): Do changes in health insurance coverage explain interstate variation in emergency department utilization?, *World Medical & Health Policy*, 8(1): 58-73
11. Regidor, E., Martinez, D., Calle, M., Astasio, P., Ortega, P., Dominguez, V. (2008): “Socioeconomic patterns in the use of public and private health services and equity in health care”, *BMC Health Services Research*, 8(183), 1-9.
12. Majo, MC and Van Soest, A (2012): “Income and health care utilization among the 50 in Europe and the US”, *Applied Econometrics*, 48: 3-41
13. Devaux, M, [et al]. (2011), “Exploring the Relationship Between Education and Obesity”, *OECD Journal: Economic Studies*, Vol. 1
14. Rodrigues, R; Ilinca, S and Schmidt, A (2017): “Income-rich and wealth-poor? The impact of measures of socio-economic status in the analysis of the distribution of long-term care use among older people”, *Health Economics*, in press, 1-10
15. Blazquez, C; Cantarero, D and Pascual, M (2017): “What Does It Drive the Relationship Between Suicides and Economic Conditions? New Evidence from Spain”, *Social Indicators*

16. DiNapoli, E; Cully, J; Mott, J; Hundt, N; Mignogna, J; Sansgiry, S; Yu, H; Trahan, L and Kunik, M (2015): “Mental health utilization among older Veterans with coexisting depression and dementia”. *SAGE Open Medicine*, 8(3).
17. Hewitt, J [et al] (2016): “Prevalence of multimorbidity and its association with outcomes in older emergency general surgical patients: an observational study”. *BMJ open*; 6(3): e010126
18. Fortin, M [et al] (2005): “Prevalence of multimorbidity among adults seen in family practice”. *Ann Fam Med*; 3: 223–28.
19. Glynn, L; Valderas, J; Healy, P; Burke, E; Newell, J; Gillespie, P and Murphy, A (2011): The prevalence of multimorbidity in primary care and its effect on health care utilization and cost. *Family Practice*; 28:516–523
20. European Commission (2014): “Communication from the Commission: On effective, accessible and resilient health systems” Available at: [https://ec.europa.eu/health/sites/health/files/systems\\_performance\\_assessment/docs/com2014\\_215\\_final\\_en.pdf](https://ec.europa.eu/health/sites/health/files/systems_performance_assessment/docs/com2014_215_final_en.pdf)
21. Riphahn, R; Wambach, A and Million, A (2003): “Incentive effects in the demand for health care: a bivariate panel count data estimation”, *Journal of Applied Econometrics*, 18(4): 387-405
22. Clavero-Barranquero, A., González-Álvarez, M.L. (2005): “Una revisión de modelos econométricos aplicados al análisis de demanda y utilización de servicios sanitarios”, *Hacienda Pública Española*, 173, 129-162.
23. Greene, W. (2012): “Econometric Analysis” 7th edition, Pearson Education.
24. Alcaide, M (2015): “Modelo de regresión binomial negativa” J. García de las Heras (Tut). Universidad de Sevilla
25. Jones, A.M.; Rice, N.; D'Uva, T.B. and Balia, S. (2013): *Applied health economics*. 2nd Edition, Routledge.
26. Börsch-Supan, A., M. Brandt, C. Hunkler, T. Kneip, J. Korbmacher, F. Malter, B. Schaan, S. Stuck, S. Zuber (2013). “Data Resource Profile: The Survey of Health, Aging and Retirement in Europe (SHARE)”, *International Journal of Epidemiology*, 42(4): 992-1001.
27. Börsch-Supan, A. (2017). *Survey of Health, Aging and Retirement in Europe (SHARE) Wave 6*. Release version: 6.0.0. SHARE-ERIC. Data set.
28. CDC-Center for Disease Control and Prevention (2015). Available at: [https://www.cdc.gov/healthyweight/assessing/bmi/adult\\_bmi/index.html](https://www.cdc.gov/healthyweight/assessing/bmi/adult_bmi/index.html)
29. Webber, L; Divajeva, D; Marsh, T; McPherson, K; Brown, M, Galea, G and Breda, J (2014): “The future burden of obesity-related diseases in the 53 WHO European-Region countries and the impact of effective interventions: a modelling study”. *BMJ Open* 2014; 4:e004787.
30. Sahoo, K; Sahoo, B; Kumar, A; Yasin, N; Kumar, R and Singh, A (2015): “Childhood obesity: causes and consequences”, *Journal of Family Medicine and Primary Care*, 4(2): 187-192.
31. Uerlich, M; Yumuk, V; Finer, N; Basdevant, A and Visscher, T (2016): “ Obesity management

- in Europe: Current status and objectives for the future”. *Obes Facts*; 9:273-283
32. Doherty, E; Queally, M; Cullinan, J and Gillespie, P (2017): “The Impact of Childhood Overweight and Obesity on Healthcare Utilization”, *Economics and Human Biology*, 27: 84.92.
  33. Crimmins, E; Kim, J and Solé-Auró, A (2010): “Gender differences in health: results from SHARE, ELSA and HRS”, *European Journal of Public Health*, 21(1). 81–91
  34. EUROSTAT (2017). Luxemburgo: Eurostat. Available at: <http://ec.europa.eu/eurostat>
  35. Ladwig, K; Marten-Mittaf, B; Formanek, B and Dammann, G (2000): “Gender differences of symptom reporting and medical health care utilization in the German population”, *European Journal of Epidemiology*, 16(6): 511-518
  36. Glaesmer, H; Brähler, E; Martin, A; Mewes, R and Rief, W (2012): “Gender differences in healthcare utilization: the mediating effect of utilization propensity”, *Journal of Applied Social Psychology*, 42(5): 1266-1279
  37. Kraut, A., Mustard, C., Walld, R., Tate, R. (2000): “Unemployment and health care utilization”, *Scandinavian Journal of Work Environment and Health*, 26(2): 169-77.
  38. Lehnert, T. (2011): “Health Care Utilization and Costs of Elderly Persons with Multiple Chronic Conditions”, *Medical care research and review* (1077-5587), 68 (4):. 387.
  39. European Parliament (1998): “Healthcare systems in the EU. A comparative study”. Available at: [http://www.europarl.europa.eu/workingpapers/saco/pdf/101\\_en.pdf](http://www.europarl.europa.eu/workingpapers/saco/pdf/101_en.pdf)
  40. Sagan A; Panteli D; Borkowski W, Dmowski M, Domański F, Czyżewski M, Goryński P, Karpacka D, Kiersztyn E, Kowalska I, Księżak M, Kuszewski K, Leśniewska A, Lipska I, Maciąg R, Madowicz J, Mądra A, Marek M, Mokrzycka A, Poznański D, Sobczak A, Sowada C, Świderek M, Terka A, Trzeciak P, Wiktorzak K, Włodarczyk C, Wojtyniak B, Wrześniewska-Wal I, Zelwianańska D, Busse R.(2011): “ Poland: Health system review”. *Health Systems in Transition*, 13(8): 1–193
  41. Kinkorova, J and Topolcan, O (2012): “Overview of healthcare system in the Czech Republic” *EPMA J.* 3 (1):4
  42. Lai T, Habicht T, Kahur K, Reinap M, Kiivet R, van Ginneken E.(2013): “Estonia: health system review”. *Health Systems in Transition*, 15(6): 1–196.
  43. Džakula A, Sagan A, Pavić N, Lončarek K and Sekelj-Kauzlarić K. (2014): “Croatia: Health system review”. *Health Systems in Transition*, 16(3): 1–162.
  44. Alexa J, Rečka L, Votápková J, van Ginneken E, Spranger A, Wittenbecher F. (2015): “Czech Republic: Health system review”. *Health Systems in Transition*, 17(1):1–165.
  45. OECD Health Systems Characteristics Survey 2016. Available at: <https://qdd.oecd.org/subject.aspx?Subject=hsc>

Country	Acronym	System
Austria	AT	Social security. 99% of population is covered. Financed by sickness funds (59%), taxation (24%), private insurance (7.5%) and co-payments
Belgium	BE	Social security. Entire population is covered. Financed by social health insurance, general taxation and out-of-pocket payments
Croatia	CRO	Mandatory Health Insurance. Universal coverage. Financed by compulsory contributions (75%) and state budget (15%) and debt.
Czech Republic	CR	Statutory Health Insurance. Compulsory membership. Financed by wage-based contributions,
Denmark	DE	National Health Service. Entire population is covered. Financed by general taxation
Estonia	ES	Estonian health system. Universal access. Financed by social payroll tax and public budget.
France	FR	National Health Insurance. Entire population is covered. Mix between private and public. Financed by employees' payroll taxes, co-payments
Germany	GE	Compulsory social insurance. Financed by compulsory and voluntary contributions to statutory health insurance (60%), general taxation (21%) and other payments
Greece	GR	Compulsory social insurance. Financed by general taxation and social insurance.
Italy	IT	National Health Service. Universal access. Financed by social insurance (40%), general taxation (35%) and co-payments.
Luxembourg	LU	Assurance Maladie. Financed by contributions to social sickness funds 50-50 employer and employee
Poland	PL	Social Health Insurance. Universal coverage. Financed by SHI contributions
Portugal	PO	National Health System. Universal coverage. Financed by general taxation (62%) and social insurance
Slovenia	SL	Health Insurance Institute of Slovenia. Universal coverage
Spain	SP	National Health System. Universal coverage. Financed by general taxation (80%), work-related contributions (18%) and co-payments.
Sweden	SW	National Insurance Scheme. Universal coverage. Financed by income taxes (65%) the rest state funds, subsidies and private insurance

**Table 1. Health care models in the European Union (Countries)**

Source: Authors' elaboration from [39-45]: European Parliament (1998), Sagan et al (2011), Kinkorova (2012), Lai et al (2013), Dzakula et al (2014), Alexa et al (2015), and OECD Health Systems Characteristics Survey (2016)

**Table 2. Variables and summary of statistics**

<i>Variable</i>	<b>Mean</b>	<b>Stand. Dev</b>	<b>Description</b>
<b>Dependent Variables</b>			
<i>GP</i>	6.7	9.136	Number of visits to the General Practitioner in 365 days
<i>HS</i>	1.781	9.1667	Hospital Stays: Number of Nights at Hospital in 365 days
<b>Independent Variables</b>			
<i>Age</i>	67.865	10.059	Age (years)
<i>Female</i>	0.560	0.496	1 if female, 0 otherwise
<i>Single</i>	0.291	0.454	1 if single, 0 otherwise
<i>InLabFor</i>	0.272	0.445	1 if in labor force, 0 otherwise
<i>SupInsu</i>	0.347	0.476	1 if has a supplementary insurance, 0 otherwise
<i>PriEduc</i>	0.367	0.481	1 if Primary School, 0 otherwise
<i>SecEduc</i>	0.538	0.498	1 if Secondary School, 0 otherwise
<i>TerEduc</i>	0.095	0.293	1 if Tertiary Education, 0 otherwise
<i>ADL</i>	0.280	0.939	Number of Limitations in Activities of Daily Living
<i>NCD</i>	1.896	1.623	Number of Chronic Diseases
<i>SPGH</i>	0.591	0.492	1 if Self-Assessed (Good Health), 0 otherwise
<i>UW</i>	0.011	0.106	1 if UnderWeight, 0 otherwise
<i>NW</i>	0.344	0.475	1 if NormalWeight, 0 otherwise
<i>OW</i>	0.418	0.493	1 if OverWeight, 0 otherwise
<i>OB</i>	0.226	0.418	1 if Obese, 0 otherwise
<i>Smoke</i>	0.448	0.497	1 if ever smoked, 0 otherwise

Source: Authors' calculation based on SHARE 6th wave (N=62,554).

**Table 3. Negative Binomial estimates. European Union. GP as dependent variable**

<i>Variable</i>	<b>50-65 years</b>	<b>65-75 years</b>	<b>75-85 years</b>	<b>+85 years</b>
<i>Female</i>	0.139 *** (0.016)	0.015 (0.017)	0.002 (0.024)	-0.077 (0.051)
<i>Single</i>	0.059 *** (0.019)	-0.025 (0.018)	-0.034 (0.023)	0.059 (0.045)
<i>SupInsu</i>	0.145 *** (0.016)	0.087 *** (0.017)	0.025 (0.022)	0.089 ** (0.040)
<i>SecEduc</i>	0.029 (0.018)	-0.001 (0.017)	0.046 ** (0.021)	0.035 (0.040)
<i>TerEduc</i>	0.115 *** (0.028)	0.054 * (0.030)	0.097 ** (0.044)	0.208 ** (0.088)
<i>ADL</i>	0.137 *** (0.015)	0.087 *** (0.012)	0.098 *** (0.011)	0.082 *** (0.012)
<i>NCD</i>	0.257 *** (0.006)	0.184 *** (0.006)	0.141 *** (0.006)	0.115 *** (0.011)
<i>SPGH</i>	-0.511 *** (0.019)	-0.439 *** (0.018)	-0.295 *** (0.023)	-0.226 *** (0.042)
<i>UW</i>	0.164 * (0.090)	0.151 * (0.087)	0.136 (0.091)	0.054 (0.121)
<i>OW</i>	-0.054 *** (0.018)	0.044 ** (0.019)	0.022 (0.024)	-0.031 (0.041)
<i>OB</i>	-0.011 (0.021)	0.058 *** (0.021)	0.006 (0.027)	-0.028 (0.055)
<i>Smoke</i>	0.002 (0.015)	0.015 (0.016)	0.045 * (0.023)	0.078 * (0.046)
<i>constant</i>	1.463 *** (0.032)	1.620 *** (0.029)	1.728 *** (0.034)	1.769 *** (0.062)
<i>alpha</i>	0.775 (0.012)	0.650 (0.011)	0.677 (0.013)	0.747 (0.025)
<i>N</i>	26,976	21,063	12,436	3,305

Notes: Standard deviations are in parentheses. \*\*\*, \*\*, \*, are the significance at level 1,5 and 10% respectively

**Table 4. Negative Binomial estimates. European Countries. GP as dependent variable**

<i>Variable</i>	<b>AT</b>	<b>BE</b>	<b>CRO</b>	<b>CR</b>	<b>DE</b>	<b>ES</b>	<b>FR</b>	<b>GE</b>
<i>Age</i>	0.041 *** (0.004)	-0.093 *** (0.021)	0.043 *** (0.005)	0.047 *** (0.003)	0.034 *** (0.005)	0.045 *** (0.003)	0.043 *** (0.004)	-0.054 ** (0.024)
<i>Age2</i>	-0.001 *** (0.000)	0.001 *** (0.000)	-0.001 *** (0.000)	-0.003 *** (0.000)	-0.001 *** (0.000)	-0.004 *** (0.000)	-0.001 *** (0.000)	0.001 ** (0.000)
<i>Female</i>	0.066 (0.041)	0.113 *** (0.034)	0.084 (0.054)	0.036 (0.029)	0.049 (0.042)	0.045 (0.038)	0.059 * (0.034)	0.045 (0.034)
<i>Single</i>	0.017 (0.044)	0.014 (0.033)	0.024 (0.081)	-0.022 (0.031)	0.033 (0.043)	-0.044 (0.036)	0.001 (0.037)	0.019 (0.041)
<i>InLabFor</i>	0.003 (0.064)	-0.312 *** (0.049)	-0.039 (0.071)	-0.088 ** (0.042)	-0.031 (0.059)	-0.074 * (0.044)	-0.003 (0.049)	-0.179 *** (0.051)
<i>SupInsu</i>	0.137 *** (0.047)	0.174 *** (0.039)	0.326 *** (0.326)	-0.012 (0.058)	0.006 (0.043)	0.111 (0.085)	0.075 (0.109)	0.093 ** (0.038)
<i>SecEduc</i>	0.046 (0.042)	-0.078 * (0.042)	-0.094 * (0.055)	0.076 ** (0.037)	0.117 * (0.063)	0.014 (0.042)	0.049 (0.037)	0.076 (0.053)
<i>TerEduc</i>	0.039 (0.069)	-0.046 (0.057)	-0.109 (0.142)	0.124 ** (0.049)	0.147 * (0.086)	0.213 *** (0.077)	0.069 (0.056)	0.146 ** (0.071)
<i>ADL</i>	0.126 *** (0.025)	0.124 *** (0.024)	0.030 (0.032)	0.069 *** (0.020)	0.095 ** (0.038)	0.093 *** (0.016)	0.108 *** (0.020)	0.073 *** (0.021)
<i>NCD</i>	0.187 *** (0.013)	0.155 *** (0.010)	0.306 *** (0.018)	0.188 *** (0.010)	0.276 *** (0.276)	0.204 *** (0.011)	0.136 *** (0.012)	0.165 *** (0.012)
<i>SPGH</i>	-0.332 *** (0.043)	-0.521 *** (0.038)	-0.397 *** (0.054)	-0.343 *** (0.031)	-0.571 *** (0.056)	-0.496 *** (0.047)	-0.483 *** (0.045)	-0.533 *** (0.039)
<i>UW</i>	0.445 ** (0.205)	0.093 (0.116)	0.212 (0.364)	-0.355 ** (0.154)	0.194 (0.189)	-0.029 (0.122)	-0.001 (0.096)	-0.148 (0.147)
<i>OW</i>	-0.039 (0.047)	0.085 ** (0.036)	-0.038 (0.063)	-0.001 (0.036)	0.076 (0.047)	-0.027 (0.043)	0.010 (0.039)	0.052 (0.037)
<i>OB</i>	0.029 (0.055)	0.068 (0.044)	0.026 (0.071)	-0.018 (0.039)	0.065 (0.057)	0.069 (0.047)	0.043 (0.043)	0.061 (0.049)
<i>Smoke</i>	0.024 (0.041)	0.075 ** (0.033)	-0.062 (0.055)	-0.005 (0.028)	-0.034 (0.042)	0.031 (0.036)	0.058 * (0.033)	-0.002 (0.033)
<i>alpha</i>	0.688 (0.029)	0.623 (0.020)	0.979 (0.044)	0.470 (0.017)	0.705 (0.030)	0.753 (0.027)	0.399 (0.024)	0.587
<i>N</i>	3,266	5,370	2294	4,612	3,499	5,356	3,718	4,233

Notes: Standard deviations are in parentheses. \*\*\*, \*\*, \*, are the significance at level 1,5 and 10% respectively. Table 1: what countries correspond to each acronym

**Table 4. Negative Binomial estimates. European Countries. GP as dependent variable (continuation)**

<i>Variable</i>	<b>GR</b>	<b>IT</b>	<b>LU</b>	<b>PL</b>	<b>PO</b>	<b>SL</b>	<b>SP</b>	<b>SW</b>
<i>Age</i>	0.025 *** (0.003)	0.053 *** (0.003)	0.055 *** (0.005)	0.051 *** (0.005)	0.027 *** (0.006)	0.051 *** (0.004)	0.045 *** (0.003)	0.037 *** (0.005)
<i>Age2</i>	-0.001 *** (0.000)	-0.001 *** (0.000)	-0.001 *** (0.000)	-0.003 *** (0.000)	-0.0001 (0.000)	-0.004 *** (0.000)	-0.003 *** (0.000)	-0.001 *** (0.000)
<i>Female</i>	0.161 *** (0.041)	0.096 *** (0.037)	0.153 ** (0.063)	0.099 * (0.055)	-0.005 (0.069)	-0.010 (0.043)	-0.043 (0.038)	-0.087 * (0.052)
<i>Single</i>	0.047 (0.043)	0.011 (0.041)	0.152 ** (0.076)	-0.125 ** (0.059)	-0.109 (0.074)	-0.027 (0.050)	-0.022 (0.042)	0.028 (0.057)
<i>InLabFor</i>	-0.232 *** (0.050)	-0.225 *** (0.048)	0.055 (0.084)	-0.337 *** (0.072)	0.009 (0.073)	0.092 (0.063)	-0.210 *** (0.047)	-0.143 ** (0.068)
<i>SupInsu</i>	0.144 * (0.083)	0.003 (0.065)	0.204 *** (0.072)	-0.035 (0.089)	0.133 * (0.069)	0.013 (0.053)	0.176 *** (0.052)	-0.106 (0.066)
<i>SecEduc</i>	-0.009 (0.039)	-0.008 (0.038)	0.023 (0.062)	0.124 ** (0.058)	0.089 (0.074)	-0.038 (0.042)	-0.048 (0.035)	0.052 (0.061)
<i>TerEduc</i>	0.034 (0.082)	-0.035 (0.069)	0.036 (0.092)	-0.043 (0.102)	0.270 (0.191)	0.058 (0.089)	0.001 (0.064)	0.209 ** (0.093)
<i>ADL</i>	0.100 *** (0.029)	0.088 *** (0.019)	0.149 *** (0.041)	-0.035 (0.031)	0.058 * (0.032)	0.014 (0.024)	0.081 *** (0.017)	0.083 ** (0.038)
<i>NCD</i>	0.263 *** (0.014)	0.204 *** (0.014)	0.151 *** (0.019)	0.212 *** (0.019)	0.107 *** (0.019)	0.174 *** (0.012)	0.165 *** (0.012)	0.218 *** (0.019)
<i>SPGH</i>	-0.383 *** (0.043)	-0.433 *** (0.041)	-0.531 *** (0.073)	-0.374 *** (0.057)	-0.434 *** (0.074)	-0.517 *** (0.044)	-0.554 *** (0.037)	-0.573 *** (0.062)
<i>UW</i>	0.078 (0.247)	0.215 (0.180)	0.031 (0.219)	0.203 (0.247)	0.237 (0.393)	0.189 (0.252)	0.214 (0.146)	-0.053 (0.229)
<i>OW</i>	0.046 (0.247)	-0.012 (0.037)	-0.110 (0.074)	0.096 (0.066)	-0.136 ** (0.069)	-0.106 ** (0.052)	-0.047 (0.037)	0.018 (0.058)
<i>OB</i>	0.063 (0.047)	0.023 (0.048)	-0.132 * (0.073)	0.120 * (0.063)	0.057 (0.086)	-0.088 (0.058)	0.097 ** (0.045)	0.024 (0.074)
<i>Smoke</i>	0.087 ** (0.039)	0.021 (0.035)	0.124 * (0.061)	-0.045 (0.053)	-0.029 (0.070)	0.029 (0.042)	0.024 (0.038)	0.074 (0.051)
<i>alpha</i>	0.871 (0.031)	0.799 (0.025)	0.600 (0.059)	0.664 (0.039)	0.574 (0.039)	0.677 (0.026)	0.561 (0.021)	0.975 (0.041)
<i>N</i>	4624	4,944	1,472	1,639	1,481	4,007	4,942	3,708

Notes: Standard deviations are in parentheses. \*\*\*, \*\*, \* are the significance at level 1,5 and 10% respectively. Table 1: what countries correspond to each acronym



**Table 5. Zero-inflated negative binomial estimates. European Union.**  
**HS as dependent variable**

<i>Variable</i>	<b>50-65 years</b>	<b>65-75 years</b>	<b>75-85 years</b>	<b>+85 years</b>
<i>Female</i>	-0.224 *** (0.071)	-0.178*** (0.066)	-0.297 *** (0.075)	0.158 (0.104)
<i>Single</i>	0.413 *** (0.088)	0.207 *** (0.072)	0.209 *** (0.081)	0.123 (0.113)
<i>SupInsu</i>	0.169 ** (0.070)	0.193 ** (0.078)	0.064 (0.071)	1.674 *** (0.161)
<i>SecEduc</i>	0.077 (0.088)	0.252 *** (0.068)	0.129 * (0.072)	-0.241 (0.194)
<i>TerEduc</i>	0.177 (0.133)	0.318 *** (0.124)	-0.008 (0.143)	0.128 *** (0.024)
<i>ADL</i>	0.249 *** (0.041)	0.381 *** (0.039)	0.205 *** (0.028)	0.249 (0.270)
<i>NCD</i>	0.355 *** (0.025)	0.258 *** (0.018)	0.059 *** (0.019)	-0.612 *** (0.141)
<i>SPGH</i>	-1.282 *** (0.076)	-1.246*** (0.066)	-0.769 *** (0.087)	0.378 *** (0.109)
<i>UW</i>	0.622 * (0.333)	0.912 *** (0.311)	0.238 (0.252)	0.110 (0.111)
<i>OW</i>	-0.237 *** (0.087)	0.028 (0.075)	-0.127 (0.082)	0.008 (0.152)
<i>OB</i>	-0.082 (0.096)	0.015 (0.077)	-0.086 (0.088)	0.046 * (0.026)
<i>Smoke</i>	0.237 *** (0.069)	0.072 (0.060)	0.031 (0.071)	0.136 (0.126)
<i>inflate cons</i>	-16.535 *** (0.282)	-17.205 (0.487)	0.921 *** (0.105)	1.538 (0.173)
<i>alpha</i>	22.119 (0.559)	16.971 (0.393)	2.102 (0.220)	21.21 (2.200)
<i>N</i>	(N=26,976)	(N=21,063)	(N= 12,436)	(N= 3,305)

Notes: Standard deviations are in parentheses. \*\*\*, \*\*, \*, are the significance at level 1,5 and 10% respectively.

**Table 6. Zero-inflated negative binomial estimates. European countries. HS as dependent variable**

<i>Variable</i>	<b>AT</b>	<b>BE</b>	<b>CRO</b>	<b>CR</b>	<b>DE</b>	<b>ES</b>	<b>FR</b>	<b>GE</b>
<i>Age</i>	0.041 *** (0.09)	0.002 (0.014)	0.060 *** (0.019)	0.043 *** (0.008)	0.002 (0.017)	0.059 *** (0.015)	-0.017 (0.018)	0.055 *** (0.009)
<i>Age2</i>	-0.001 ** (0.0001)	0.001 (0.000)	-0.001 *** (0.000)	-0.001 ** (0.000)	-0.001 (0.000)	-0.001 *** (0.000)	0.001 (0.000)	-0.001 *** (0.000)
<i>Female</i>	0.026 (0.110)	-0.286 ** (0.141)	-0.153 (0.184)	0.058 (0.105)	0.023 (0.173)	-0.342 ** (0.144)	-0.015 (0.180)	-0.158 * (0.091)
<i>Single</i>	0.082 (0.107)	0.127 (0.148)	-0.129 (0.206)	0.202 * (0.107)	0.212 (0.162)	0.461 *** (0.141)	0.016 (0.181)	0.462 *** (0.106)
<i>InLabFor</i>	-0.214 (0.186)	-0.732 *** (0.197)	-0.293 (0.267)	-0.260 (0.160)	-0.815 *** (0.235)	-0.749 *** (0.202)	-0.779 *** (0.256)	-0.373 *** (0.125)
<i>SupInsu</i>	0.461 *** (0.145)	-0.303 (0.188)	0.714 ** (0.0358)	-0.037 (0.214)	-0.294 * (0.173)	0.048 (0.219)	0.147 (0.396)	0.064 (0.093)
<i>SecEduc</i>	-0.001 (0.112)	0.346 ** (0.137)	-0.143 (0.183)	0.077 (0.118)	0.289 (0.199)	-0.021 (0.141)	0.345 (0.227)	0.102 (0.135)
<i>TerEduc</i>	-0.183 (0.214)	0.128 (0.180)	-0.209 (0.609)	0.031 (0.173)	1.347 *** (0.327)	0.438 (0.267)	0.379 (0.322)	-0.226 (0.184)
<i>ADL</i>	0.119 *** (0.045)	0.289 *** (0.072)	-0.002 (0.054)	0.177 *** (0.033)	0.316 ** (0.124)	0.275 *** (0.048)	0.294 *** (0.076)	0.105 *** (0.035)
<i>NCD</i>	0.184 *** (0.032)	0.163 *** (0.042)	0.129 ** (0.060)	0.089 *** (0.028)	0.344 *** (0.059)	0.169 ** (0.066)	0.143 ** (0.060)	0.156 *** (0.034)
<i>SPGH</i>	-0.771 *** (0.120)	-1.284 *** (0.153)	-1.269 *** (0.296)	-0.643 *** (0.111)	-1.136 *** (0.196)	-1.025 *** (0.223)	-1.715 *** (0.191)	-1.019 *** (0.123)
<i>UW</i>	-0.016 (0.299)	-0.826 ** (0.371)	0.099 (0.440)	0.507 (0.681)	0.741 (0.789)	-0.489 (0.339)	0.654 (0.548)	0.759 *** (0.227)
<i>OW</i>	-0.036 (0.119)	-0.106 (0.154)	0.207 (0.202)	-0.146 (0.124)	-0.207 (0.199)	-0.353 ** (0.169)	0.206 (0.199)	0.050 (0.105)
<i>OB</i>	-0.029 (0.132)	0.172 (0.178)	-0.037 (0.241)	-0.215 * (0.125)	-0.4431 * (0.252)	-0.386 ** (0.157)	-0.013 (0.241)	0.162 (0.121)
<i>Smoke</i>	0.194 * (0.105)	0.268 ** (0.131)	-0.326 ** (0.181)	0.162 * (0.093)	0.007 (0.169)	0.051 (0.124)	0.449 *** (0.171)	0.073 (0.093)
<i>infl cons</i>	0.671 *** (0.132)	-14.436 *** (0.529)	1.442 *** (0.266)	1.146 *** (0.067)	-14.864 *** (0.268)	0.0886 *** (0.296)	-13.074 *** (0.534)	0.909 *** (0.094)
<i>alpha</i>	1.852 (0.339)	14.529 (0.694)	2.323 (0.927)	1.430 (0.181)	16.154 (1.134)	3.567 (1.287)	17.564 (0.063)	1.663 (0.237)
<i>N</i>	3,266	5,370	2,294	4,612	3,499	5,356	3,718	4,233

Notes: Standard deviations are in parentheses. \*\*\*, \*\*, \*, are the significance at level 1, 5 and 10% respectively. Table 1: what countries correspond to each acronym

**Table 6. Zero-inflated negative binomial estimates. European countries. HS as dependent variable (continuation)**

<i>Variable</i>	<b>GR</b>	<b>IT</b>	<b>LU</b>	<b>PL</b>	<b>PO</b>	<b>SL</b>	<b>SP</b>	<b>SW</b>
<i>Age</i>	-0.425 *** (0.114)	0.014 (0.029)	0.061 *** (0.015)	0.091 *** (0.11)	-0.820 *** (0.251)	-0.223 * (0.126)	0.023 * (0.013)	-0.041 *** (0.014)
<i>Age2</i>	0.003 *** (0.001)	-0.001 (0.000)	-0.004 *** (0.000)	-0.001 *** (0.000)	0.006 *** (0.002)	0.002 * (0.001)	-0.001 (0.000)	0.001 *** (0.000)
<i>Female</i>	-0.773 *** (0.207)	-0.484 *** (0.167)	-0.143 (0.188)	-0.036 (0.172)	-0.891 *** (0.276)	-0.228 (0.139)	-1.003 *** (0.175)	-0.196 (0.149)
<i>Single</i>	0.285 (0.211)	0.169 (0.185)	0.462 ** (0.195)	0.219 (0.185)	0.483 (0.331)	0.173 (0.45)	0.548 *** (0.204)	0.160 (0.176)
<i>InLabFor</i>	-0.307 (0.246)	-0.301 (0.232)	-0.689 ** (0.285)	-0.512 * (0.279)	-0.250 (0.364)	-0.729 *** (0.261)	-1.006 *** (0.231)	-0.469 ** (0.209)
<i>SupInsu</i>	0.413 (0.378)	-0.139 (0.352)	0.368 ** (0.186)	0.402 (0.379)	-0.015 (0.279)	-0.149 (0.149)	0.682 ** (0.268)	0.113 (0.223)
<i>SecEduc</i>	-0.818 *** (0.210)	0.268 (0.193)	-0.213 (0.166)	0.036 (0.158)	-0.337 (0.326)	-0.233 (0.142)	-0.109 (0.167)	0.084 (0.166)
<i>TerEduc</i>	-0.673 * (0.363)	0.149 (0.284)	-0.582 ** (0.269)	0.373 (0.486)	-1.239 *** (0.459)	-0.582 * (0.327)	-0.919 *** (0.286)	0.178 (0.250)
<i>ADL</i>	0.464 *** (0.095)	0.271 *** (0.091)	0.341 *** (0.077)	-0.022 (0.041)	0.398 *** (0.111)	0.154 *** (0.059)	0.379 *** (0.089)	0.188 ** (0.085)
<i>NCD</i>	0.307 *** (0.064)	0.266 *** (0.072)	0.074 * (0.045)	0.065 (0.042)	0.191 *** (0.071)	0.183 *** (0.052)	0.197 *** (0.053)	0.234 *** (0.049)
<i>SPGH</i>	-1.98 *** (0.192)	-1.825 *** (0.185)	-0.829 *** (0.178)	-0.973 *** (0.198)	-1.094 *** (0.320)	-0.918 *** (0.165)	-1.733 *** (0.167)	-0.924 *** (0.153)
<i>UW</i>	1.150 (0.949)	1.173 (0.845)	1.285 *** (0.484)	0.636 (0.605)	-0.693 (0.625)	0.391 (0.414)	0.186 (0.571)	0.504 (0.528)
<i>OW</i>	0.136 (0.207)	-0.181 (0.178)	-0.366 * (0.202)	-0.380 ** (0.181)	-0.916 *** (0.283)	-0.344 ** (0.156)	-0.545 *** (0.192)	-0.008 (0.174)
<i>OB</i>	0.520 * (0.276)	-0.242 (0.189)	0.074 (0.196)	-0.291 * (0.174)	-0.931 ** (0.384)	-0.332 * (0.189)	-0.239 (0.211)	0.288 (0.202)
<i>Smoke</i>	0.081 (0.195)	0.034 (0.148)	0.172 (0.178)	-0.059 (0.156)	-0.262 (0.278)	0.184 (0.129)	-0.098 (0.169)	0.113 (0.148)
<i>infl cons</i>	-13.354 *** (0.383)	-0.415 (2.198)	1.217 *** (0.158)	1.239 *** (0.093)	-12.636 *** (1.416)	0.951 *** (0.232)	-14.716 *** (0.368)	-13.059 *** (2.236)
<i>alpha</i>	23.936 (0.073)	11.775 (12.363)	1.789 (0.440)	1.354 (0.211)	20.623 (1.895)	3.019 (0.903)	19.534 (1.069)	14.632 (0.089)
<i>N</i>	4,624	4,944	1,472	1,639	1,481	4,007	4,942	3,708

Notes: Standard deviations are in parentheses. \*\*\*, \*\*, \* are the significance at level 1, 5 and 10% respectively. Table 1: what countries correspond to each acronym